

Cambridge International AS Level

MATHEMATICS

Paper 2 Pure Mathematics 2

MARK SCHEME

Maximum Mark: 50

Published	
Published	

Students did not sit exam papers in the June 2020 series due to the Covid-19 global pandemic.

This mark scheme is published to support teachers and students and should be read together with the question paper. It shows the requirements of the exam. The answer column of the mark scheme shows the proposed basis on which Examiners would award marks for this exam. Where appropriate, this column also provides the most likely acceptable alternative responses expected from students. Examiners usually review the mark scheme after they have seen student responses and update the mark scheme if appropriate. In the June series, Examiners were unable to consider the acceptability of alternative responses, as there were no student responses to consider.

Mark schemes should usually be read together with the Principal Examiner Report for Teachers. However, because students did not sit exam papers, there is no Principal Examiner Report for Teachers for the June 2020 series.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the June 2020 series for most Cambridge IGCSE™ and Cambridge International A & AS Level components, and some Cambridge O Level components.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

© UCLES 2020 Page 2 of 12

Mathematics-Specific Marking Principles 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing. 2 Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected. 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points. 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw). 5 Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread. 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

© UCLES 2020 Page 3 of 12

Mark Scheme Notes

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- **B** Mark for a correct result or statement independent of method marks.
- DM or DB When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
 - FT Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.

© UCLES 2020 Page 4 of 12

Abbreviations

AEF/OE Any Equivalent Form (of answer is equally acceptable) / Or Equivalent

AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)

CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)

CWO Correct Working Only

ISW Ignore Subsequent Working

SOI Seen Or Implied

SC Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the

light of a particular circumstance)

WWW Without Wrong Working

ARWT Answer Which Rounds To

© UCLES 2020 Page 5 of 12

Question	Answer	Marks
1	Apply logarithms to both sides and apply power law at least once	M1
	Rearrange to the form $y = \frac{3 \ln 9}{\ln 2} x$ OE	A1
	Obtain $k = 9.51$	A1
		3

Question	Answer	Marks
2	Differentiate using product rule to obtain $ae^{\frac{1}{2}x} + bxe^{\frac{1}{2}x}$	*M1
	Obtain correct $5e^{\frac{1}{2}x} + \frac{5}{2}xe^{\frac{1}{2}x}$ OE	A1
	Equate first derivative to zero and solve for <i>x</i>	DM1
	Obtain <i>x</i> -coordinate –2	A1
	Obtain y-coordinate $-10e^{-1}$	A1
		5

© UCLES 2020 Page 6 of 12

Question	Answer	Marks
3	Differentiate $\cos 3x$ to obtain $-3\sin 3x$	B1
	Differentiate $5\sin y$ to obtain $5\cos y \frac{dy}{dx}$	B1
	Obtain $-3\sin 3x + 5\cos y \frac{dy}{dx} = 0$ OE	B1
	Substitute <i>x</i> and <i>y</i> values to find value of first derivative	M1
	Obtain $\frac{3}{5}$	A1
		5

© UCLES 2020 Page 7 of 12

Question	Answer	Marks
4	State or imply equation is $\ln y = \ln A - 2p \ln x$	B1
	Equate gradient of line to $-2p$	M1
	Obtain $-2p = -2.6$ and hence $p = 1.3$	A1
	Substitute appropriate values to find ln A	M1
	Obtain $\ln A = 1.252$ and hence $A = 3.5$	A1
	Alternative method for question 4	
	State or imply equation is $\ln y = \ln A - 2p \ln x$	B1
	Substitute given coordinates to obtain 2 simultaneous equations and solve to obtain 3.5p	M1
	Obtain $3.5p = 4.55$ and hence $p = 1.3$	A1
	Substitute appropriate values to find ln A	M1
	Obtain $\ln A = 1.252$ and hence $A = 3.5$	A1
		5

© UCLES 2020 Page 8 of 12

Question	Answer	Marks
5(a)	Draw V-shaped graph with vertex on positive <i>x</i> -axis	B1
	Draw (more or less) correct graph of $y = 3x + 5$	B1
		2
5(b)	State equation $3x + 5 = -(2x - 3)$ or corresponding inequality	B1
	Attempt solution of linear equation / inequality where signs of $3x$ and $2x$ are different	M1
	State answer $x < -\frac{2}{5}$	A1
	Alternative method for question 5(b)	
	Square both sides of equation / inequality and attempt solution of 3-term quadratic equation / inequality	M1
	Obtain (eventually) only $-\frac{2}{5}$	A1
	State answer $x < -\frac{2}{5}$	A1
		3

© UCLES 2020 Page 9 of 12

Question	Answer	Marks
6(a)	Substitute $x = -3$, equate to zero and attempt solution for a	M1
	Obtain $a = 17$	A1
		2
6(b)	Divide by $x+3$ at least as far as the x term	M1
	Obtain $6x^2 - x - 1$	A1
	Conclude $(x+3)(3x+1)(2x-1)$	A1
		3
6(c)	Attempt solution of $\sin \theta = k$ where $-1 \le k \le 1$	M1
	Obtain 199.5	A1
	Obtain 340.5	A1
		3

Question	Answer	Marks
7(a)	Integrate to obtain the form $k_1 \ln(2x+1) + k_2 x^2$	*M1
	Obtain correct $2\ln(2x+1) + 4x^2$	A1
	Use limits correctly and attempt rearrangement	DM1
	Confirm $a = \sqrt{2.5 - 0.5 \ln(2a + 1)}$ AG	A1
		4

© UCLES 2020 Page 10 of 12

Question	Answer	Marks
7(b)	Consider sign of $a - \sqrt{2.5 - 0.5 \ln(2a + 1)}$ or equivalent for 1 and 2	M1
	Obtain -0.3 and 0.6 or equivalents and justify conclusion	A1
		2
7(c)	Use iteration process correctly at least once	M1
	Obtain final answer 1.358	A1
	Show sufficient iterations to 6 sf to justify answer or show a sign change in the interval [1.3575, 1.3585]	A1
		3

Question	Answer	Marks
8(a)	Use at least one of $\sin 2\theta = 2\sin \theta \cos \theta$ and $\cot \theta = \frac{\cos \theta}{\sin \theta}$	B1
	Use both and conclude $6\cos^2\theta$ AG	B1
		2
8(b)	Attempt solution of $\cos^2 \theta = \frac{5}{6}$ to find at least one value	M1
	Obtain 0.421	A1
	Obtain 2.72	A1
		3

© UCLES 2020 Page 11 of 12

Question	Answer	Marks
8(c)	Express integrand in form $a + b \cos x$	M1
	Obtain correct integrand $3 + 3\cos x$	A1
	Integrate to obtain $px + q \sin x$	*M1
	Apply limits correctly	DM1
	Obtain $\frac{3}{4}\pi + 3 - \frac{3}{\sqrt{2}}$ or exact equivalent	A1
		5

© UCLES 2020 Page 12 of 12