

## **MARK SCHEME for the May/June 2013 series**

### **0620 CHEMISTRY**

**0620/31**

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0620	31

- 1 (a) (i) contains carbon and hydrogen [1]  
**cond:** only / just [1]
- (ii) (different) boiling points [1]  
**cond:** separate [1]
- (b) bitumen-making roads / roofs / water-proofing, etc. [1]
- lubricating fraction – waxes / vaseline / grease, etc. or machinery example, e.g. (oil a) bike / hinges / reducing friction [1]
- paraffin fraction – jet fuel / (home) heating or tractors or cooking or lighting [1]
- gasoline fraction – petrol or fuel for cars / vans / trucks [1]
- [Total: 8]**
- 2 (a) 3 or III [1]
- (b) good conductor and it is a metal/has delocalised (free) electrons [1]
- (c) N or P or As or Sb [1]  
**accept Bi**
- (d)  $M_2(SO_4)_3$  [1]  
**accept:**  $Ga_2(SO_4)_3$
- (e) it would react with/dissolves in a named strong acid [1]  
it would react with/dissolves in a named alkali [1]  
it shows both basic and acid properties =1 [1]  
it reacts with both acids and bases/alkalis =1 [1]  
[**max 2**]
- [Total: 6]**

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0620	31

- 3 (a) (i) pieces have (same) surface area [1]  
 same amount / mass / quantity / volume / number of moles of carbonate [1]
- (ii) no more bubbles / carbon dioxide **or** piece disappears / dissolves [1]
- (b) experiment 1  $\text{Ca}^{2+} + \text{CO}_2 + \text{H}_2\text{O}$  [1]
- (c) (i) more concentrated **or** higher concentration (of acid) (in experiment 1) [1]  
 accept: arguments based on collision theory
- (ii) ethanoic acid is a weak acid **or** hydrochloric acid is a strong acid [1]  
 accept: stronger or weaker
- ethanoic acid less ionised / dissociated / lower / smaller concentration of hydrogen ions [1]  
 accept: less hydrogen ions and vice versa argument but not dissociation of ions
- (iii) lower temperature (particles) have less energy [1]  
 moving more slowly [1]  
 fewer collisions / lower collision rate [1]  
**or**  
 lower temperature (particles) have less energy [1]  
 fewer particles collide [1]  
 with the necessary energy to react [1]  
 note: less energy fewer successful collisions gains all 3 marks
- [Total: 10]**
- 4 (a) it is an alkane **or** hydrocarbon [1]  
 it is saturated **or** only C—C single bonds [1]  
 accept: no double bonds
- (b) molecular formula  $\text{C}_6\text{H}_{12}$  [1]  
 empirical formula  $\text{CH}_2$  [1]
- (c) correct structural formula of cyclobutane [1]

Page 4	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0620	31

(d) (i)  $C_6H_{12}$  [1]  
**accept:** a correct structural formula

(ii) same molecular formula **not:** chemical formula [1]  
different structural formulae / structures [1]

(e) add bromine (water) or (l) [1]

**cond:** (remains) brown **or** orange **or** red or yellow [1]

**cond:** changes from brown, etc. to colourless or decolourises [1]  
**not:** clear

**OR**

potassium manganate(VII) [1]

**note:** oxidation state not essential but if given must be correct or [0]

**accept:** potassium permanganate

**cond:** remains pink / purple [1]

**cond:** changes from pink to colourless (**acidic**) [1]  
**not:** clear

**cond:** change from pink to green / brown (**alkaline**)

[Total: 11]

5 (a) (i) any metal above zinc [1]  
 $Mg \rightarrow Mg^{2+} + 2e^{-}$

(ii)  $Zn + 2Ag^{+} \rightarrow Zn^{2+} + 2Ag$  [2]  
**Note:** not balanced only [1]

(iii) because they can accept or gain electrons / change into atoms or can be reduced [1]

(iv)  $Ag^{+}$  or silver [1]  
charge not essential but if given must be correct

(v)  $Ag^{+}$  and  $Cu^{2+}$  **or** silver and copper [1]  
charge not essential but if given must be correct

Page 5	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0620	31

- (b) Cu Sn Cd Zn (*i.e.* all 4 in correct order) [1]  
relates order to voltage [1]

one relevant comment from: [1]

higher reactivity metals are the negative electrode / copper is least reactive because it is the positive electrode because copper would have the lowest voltage / copper cell  $V = 0$  / the bigger the difference in reactivity, the bigger the voltage / zinc has highest voltage because it is most reactive / more reactive metals have higher voltage

[Total: 9]

- 6 (a) (i) proton or  $H^+$  acceptor [1]
- (ii) (measure) pH or (use) UI indicator [1]  
**note:** can be implied need not be explicit  
sodium hydroxide has higher pH / ammonia(aq) has lower pH [1]  
(this sentence would score 2 marks)  
**or**  
appropriate colours with UI / appropriate numerical values [1]  
ammonia is closer to green, blue-green, turquoise or lighter blue  
sodium hydroxide is darker blue / purple / violet [1]  
**or**  
measure electrical conductivity [1]  
can be implied need not be explicit  
ammonia (aq) is the poorer conductor/ sodium hydroxide is the better conductor [1]

Page 6	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0620	31

(b) any five from:

- high pressure favours lower volume side / movement to right / ammonia side, **or** high pressure increases the yield
- high pressure increases rate
- low temperature favours exothermic reaction / increases yield / favours the forward reaction
- low temperature gives low rate or vice versa
- catalyst increases rate or lowers activation energy
- 450 °C low enough to give an economic yield but with catalyst gives a fast enough rate  
note need whole concept to get this compromise temperature point [5]

(c)  $2\text{NH}_3 + \text{NaClO} \rightarrow \text{N}_2\text{H}_4 + \text{NaCl} + \text{H}_2\text{O}$  [2]  
not balanced only 1

(d) 4 hydrogen atoms 1 bonding pair each [1]  
2 nitrogen atoms with 1 bonding pair between them [1]  
one non-bonding pair on each N (need not be seen as a pair) [1]

(e) (i) pH increases [1]

(ii) oxygen needed for rusting / removes oxygen / reacts with oxygen [1]

**[Total: 15]**

7 (a) (i) add carbon / animal charcoal [1]  
filter [1]

**OR**

repeat experiment without indicator [1]  
using same quantity / volume of acid [1]

(ii) add magnesium metal / carbonate / oxide / hydroxide [1]  
to (hot) (hydrochloric) acid [1]

**cond:** until in excess **or** no more dissolves **or** reacts [1]

**cond:** filter (to remove unreacted solid) [1]

Page 7	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0620	31

- (b) number of moles of HCl =  $0.020 \times 2.20 = 0.044$  [1]  
 number of moles of LiOH = 0.044  
 concentration of LiOH =  $0.044/0.025 = 1.769$  (mol/dm<sup>3</sup>) [1]  
**accept** 1.75 to 1.77 need 2 dp  
 correct answer scores = 2

- (c) (for LiCl·2H<sub>2</sub>O)  
 mass of one mole = 78.5 [1]  
 percentage water =  $36 / 78.5 \times 100$  [1]  
 45.9 so is LiCl·2H<sub>2</sub>O [1]  
 only award the marks if you can follow the reasoning and it gives 45.9% of water

**note:** if correct option given mark this and ignore the rest of the response

**allow:** max 2 for applying a correct method to another hydrate, [1] for the method and [1] for the correct value, working essential

[Total: 10]

- 8 (a) (i) regular arrangement / repeating pattern **NOT** structure [1]  
**cond:** ions [1]  
**not** molecules / atoms
- (ii) attraction between opposite charges / electrostatic attraction [1]
- (b) delocalised / mobile / free / sea of electrons [1]  
 positive ions / cations  
**not** atoms / protons / nuclei [1]  
 attraction between these electrons and ions [1]
- (c) **giant covalent**  
 no ions [1]  
 no delocalised / free / mobile / sea of electrons **or** all electrons [1]
- ionic**  
 in ionic solid ions cannot move [1]  
 liquid ionic compound ions can move [1]
- metallic**  
 (both solid and liquid) metals have delocalised (**or** alternative term) electrons [1]

[Total: 11]