



**Cambridge International Examinations**  
Cambridge International General Certificate of Secondary Education

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 1 7 9 3 5 6 9 0 2 4 \*



**CHEMISTRY**

**0620/52**

Paper 5 Practical Test

**February/March 2016**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Practical notes are provided on pages 11 and 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

**For Examiner's Use**

**Total**

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **9** printed pages and **3** blank pages.

- 1 You are going to investigate the rate of a reaction between two solutions, **J** and **K**, and sulfuric acid at different temperatures.

**Read all the instructions below carefully before starting the experiments.**

### Instructions

You are going to carry out four experiments.

#### (a) Experiment 1

Use the large measuring cylinder to pour 50 cm<sup>3</sup> of distilled water and 40 cm<sup>3</sup> of sulfuric acid into the 250 cm<sup>3</sup> conical flask.

Use the small measuring cylinder to add 2 cm<sup>3</sup> of methyl orange and 5 cm<sup>3</sup> of solution **J** to the mixture in the conical flask. Measure the temperature of the mixture and record it in the table.

Use the small measuring cylinder to now start the reaction by adding 5 cm<sup>3</sup> of solution **K** to the conical flask and immediately start your timer and swirl the mixture.

Measure the time taken for the mixture to turn very pale yellow and record the time in the table.

Measure and record the final temperature of the mixture.

#### Experiment 2

Repeat Experiment 1 but heat the mixture in the conical flask to about 30 °C **before** adding the solution **K**. Measure and record the temperature of the mixture.

Now add 5 cm<sup>3</sup> of solution **K** to the conical flask and immediately start your timer and swirl the mixture.

Measure the time taken for the mixture to turn very pale yellow and record the time in the table.

Measure and record the final temperature of the mixture.

#### Experiment 3

Repeat Experiment 1 but heat the mixture in the conical flask to about 40 °C before adding the solution **K** to the conical flask. Take the same measurements and record the values in the table.

#### Experiment 4


Repeat Experiment 1 but heat the mixture in the conical flask to about 50 °C before adding the solution **K** to the conical flask. Take the same measurements and record the values in the table.

Work out the average temperatures to complete the table.

experiment	time taken for mixture to turn very pale yellow / s	initial temperature / °C	final temperature / °C	average temperature / °C
1				
2				
3				
4				

[4]

(b) Plot the results you have obtained on the grid and draw a smooth line graph.



[4]

(c) **From your graph**, deduce the time taken for the mixture to turn very pale yellow if Experiment 1 was repeated at an average temperature of 60 °C.  
Show clearly **on the grid** how you worked out your answer.

..... [2]

(d) (i) In which experiment was the rate of reaction greatest?

..... [1]

(ii) Explain why the rate of reaction was greatest in this experiment.

.....  
.....  
..... [2]

(e) (i) Suggest and explain the effect **on the results** of using a burette to measure the volume of solution J.

.....  
..... [2]

(ii) Suggest and explain one **other** improvement to these experiments.

.....  
..... [2]

[Total: 17]

- 2 You are provided with two solids, **L** and **M**, which are both salts.  
Carry out the following tests on the solids, recording all of your observations at each stage.

**tests on solid L**

- (a) Describe the appearance of solid **L**.

..... [1]

- (b) Add about 10 cm<sup>3</sup> of distilled water to solid **L** and shake the mixture for about one minute to dissolve.

Divide the solution into four equal portions in four test-tubes and carry out the following tests.

- (i) Add several drops of aqueous ammonia to the first portion of the solution.

Now add excess ammonia solution to the mixture and shake.

observation .....

.....

.....

..... [4]

- (ii) Add excess aqueous sodium hydroxide to the second portion of the solution.

observation .....

..... [1]

- (iii) Add about 1 cm<sup>3</sup> of dilute nitric acid to the third portion of the solution followed by aqueous silver nitrate.

observation ..... [1]

- (iv) Add about 1 cm<sup>3</sup> of dilute nitric acid to the fourth portion of the solution followed by aqueous barium nitrate.

observation ..... [1]

- (c) Identify solid **L**.

.....

..... [2]

**tests on solid M**

- (d) Use a spatula to divide solid **M** into two portions in two test-tubes.

Heat the first portion of the solid, **gently**.  
Test the gas given off with damp red litmus paper.

observation .....

.....

..... [3]

- (e) Dissolve the second portion of the solid in about 6 cm<sup>3</sup> of water. Pour half of the solution into a boiling tube.

- (i) Add about 1 cm<sup>3</sup> of aqueous sodium hydroxide to the solution in the boiling tube and heat the mixture gently. Test the gas given off.

observation ..... [1]

- (ii) Add about 1 cm<sup>3</sup> of dilute nitric acid to the remaining solution in the test-tube followed by aqueous silver nitrate.

observation ..... [1]

- (f) Identify solid **M**.

.....

..... [2]

[Total: 17]











**NOTES FOR USE IN QUALITATIVE ANALYSIS****Test for anions**

anion	test	test result
carbonate ( $\text{CO}_3^{2-}$ )	add dilute acid	effervescence, carbon dioxide produced
chloride ( $\text{Cl}^-$ ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
bromide ( $\text{Br}^-$ ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	cream ppt.
iodide ( $\text{I}^-$ ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate ( $\text{NO}_3^-$ ) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate ( $\text{SO}_4^{2-}$ ) [in solution]	acidify, then add aqueous barium nitrate	white ppt.
sulfite ( $\text{SO}_3^{2-}$ )	add dilute hydrochloric acid, warm gently and test for the presence of sulfur dioxide	sulfur dioxide produced will turn acidified aqueous potassium manganate(VII) from purple to colourless

**Test for aqueous cations**

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium ( $\text{Al}^{3+}$ )	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium ( $\text{NH}_4^+$ )	ammonia produced on warming	—
calcium ( $\text{Ca}^{2+}$ )	white ppt., insoluble in excess	no ppt., or very slight white ppt.
chromium(III) ( $\text{Cr}^{3+}$ )	green ppt., soluble in excess	grey-green ppt., insoluble in excess
copper ( $\text{Cu}^{2+}$ )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess, giving a dark blue solution
iron(II) ( $\text{Fe}^{2+}$ )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) ( $\text{Fe}^{3+}$ )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc ( $\text{Zn}^{2+}$ )	white ppt., soluble in excess, giving a colourless solution	white ppt., soluble in excess, giving a colourless solution

**Test for gases**

gas	test and test results
ammonia (NH <sub>3</sub> )	turns damp, red litmus paper blue
carbon dioxide (CO <sub>2</sub> )	turns limewater milky
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper
hydrogen (H <sub>2</sub> )	'pops' with a lighted splint
oxygen (O <sub>2</sub> )	relights a glowing splint
sulfur dioxide (SO <sub>2</sub> )	turns acidified aqueous potassium manganate(VII) from purple to colourless

**Flame tests for metal ions**

metal ion	flame colour
lithium (Li <sup>+</sup> )	red
sodium (Na <sup>+</sup> )	yellow
potassium (K <sup>+</sup> )	lilac
copper(II) (Cu <sup>2+</sup> )	blue-green

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.